

Above-Ground Carbon Stock

Indicator name Above-Ground Carbon Indicator (AGCI)

Indicator unit The above-ground carbon (AGC) is expressed in Mg (megagrams or tonnes) of carbon per km². It corresponds to the carbon fraction of the oven-dry weight of the woody parts (stem, bark, branches and twigs) of all living trees, excluding stump and roots, as estimated by the GlobBiomass project (globbiomass.org) with 2010 as the reference year.

Area of interest The AGCI has been calculated in DOPA for each terrestrial and coastal protected area of size ≥ 25 km² and at country and ecoregion levels.

Related targets



[Sustainable Development Goal 13 on climate action](#)



[Sustainable Development Goal 15 on life on land](#)



[Aichi Biodiversity Target 11 on protected areas](#)



[Aichi Biodiversity Target 15 on contribution to carbon stocks](#)

Policy question There are two main policy questions to which AGCI is relevant:

- How do protected areas contribute, through the conservation of vegetation resources, to the health and productivity of the ecosystems and to the sustainability of the local communities that depend on these resources? The persistent loss of AGC can indicate a degradation of the forest vegetation canopy, and can happen through unsustainable management practices and through detrimental land use and land cover changes.
- How do protected areas contribute to carbon storage and hence to offset the impacts of fossil fuel emissions and to climate change mitigation? Forests represent one of the largest terrestrial organic carbon reservoirs, and significantly contribute to the regulation of the global carbon cycle. Changes in land use and land cover can cause AGC decreases and related carbon emissions, which are one of the largest sources of human-caused carbon emissions to the atmosphere. Protected areas may contribute to biomass and carbon retention and hence to the reduction of net emissions of greenhouse gases responsible for climate change.

Use and interpretation

Tree carbon stocks are relevant for quantifying terrestrial carbon storage and carbon sinks as well as for estimating potential emissions from land cover

changes (deforestation, reforestation, afforestation) and from biotic (pests, diseases) and abiotic (e.g. forest fires, windstorms) disturbances. Forests in particular have a key role in the global carbon cycle and are considered large and persistent carbon sinks. The CO₂ fixed by photosynthesis is one of the most important components of the carbon cycle, and forests play a determinant role in this process. Therefore, spatially explicit data and assessments of forest biomass and carbon are of paramount importance for the design and implementation of effective sustainable forest management options and forest related policies.

The AGCI provides useful information about the tree carbon stocks and condition in protected areas, which can contribute to identify potentially degraded areas, evaluate the conservation performance of protected areas, set restoration targets, and assess the contribution of protected areas to reduce net global carbon emissions.

In the assessment of AGCI, water bodies, urban areas, permanent snow/ice and bare area land cover classes mapped by the Climate Change Initiative – Land Cover map (Land Cover CCI, 2017) have been masked out for preventing distortions and potentially biased estimates that unvegetated areas, or areas with very low canopy cover, can cause in the assessment (Quegan *et al.* 2017).

Key caveats

The AGCI was obtained by converting biomass to carbon using a conversion factor of 0.5 applied to the above-ground biomass (AGB) data provided by the global terrestrial biomass map derived from Earth Observation data in the framework of the GlobBiomass project (<http://globbiomass.org>) funded by the European Space Agency (ESA). The AGB estimates in the GlobBiomass project were derived from spaceborne SAR (ALOS PALSAR, Envisat ASAR), optical (Landsat-7), LiDAR (ICESAT) satellite data and auxiliary datasets using multiple estimation procedures. Uncertainties and accuracy of these estimates, and hence of the AGCI, vary in space and time.

Overall, the GlobBiomass approach seems to be performing appropriately in estimating AGB in all biomes, as assessed over a significant set of locations with independent in situ reference data (Rozendaal *et al.*, 2017). The validation confirmed the quality of the AGB estimates and indicated reliability even in the wet tropics, which was initially considered to be beyond the capability of the EO datasets and algorithms available. The estimates, however, are not free from errors (local biases and substantial uncertainties), primarily in regions where the remote sensing data available was of limited usefulness to resolve forest structures or to be sufficiently characterized in terms of wood density and biomass expansion factors. For all biomes, AGB predictions strongly agreed with the observations in the lower biomass range. In the temperate and subtropical zone, AGB was underestimated for reference values ≥ 150 Mg/ha, while in the tropics and in the boreal ecozone the agreement between predicted and observed AGB was remarkable. We refer to the documentation of the GlobBiomass project (Quegan *et al.*, 2017; Rozendaal *et al.*, 2017) for a detailed discussion about the main strengths and limitations of the product.

Trees are the main stock of terrestrial vegetation biomass and carbon, but other vegetation types, such as shrubs or herbaceous plants can provide also significant contributions to AGB, which are not considered by the AGCI.

AGB estimates were generated in GlobBiomass project for each point on Earth for which EO data were available. However, areas with no or very low canopy cover (typically, water, urban, permanent snow and ice and bare soil) have been masked out and are not considered in the assessment.

The biomass to carbon conversion factor of 0.5 here used is a good approximation of the typical carbon content in the biomass of terrestrial vegetation, and is consistent with the approach in the Good Practice Guidance in LULUCF by the IPCC (2003) and in other related assessments (Baccini et al., 2017; Zarin et al., 2016; Achard et al., 2014; Baccini et al., 2012; Saatchi et al., 2011; Gallaun et al., 2010). There is however some variation of this biomass to carbon conversion factor for different tree species, different components of a tree or a stand and age of the stand, which may be accounted for in more detailed assessments (Ruesch and Gibbs, 2008; Thurner et al., 2014).

Because the AGCI is computed within the boundaries for each protected area with a size of at least 25 km², results will be affected by the accuracy of the available protected area boundaries (see Figure 1).

Indicator status The Above Ground Biomass map, developed by ESA's GlobBiomass project, is publicly available for download at <http://globbiomass.org/products/global-mapping/> (Santoro *et al.*, 2018) and is described in detail in Quegan *et al.* (2017). The assessment of the AGCI in protected areas is not yet published.

Available data and resources

Data available AGCI values are available for each protected area of size ≥ 25 km², and can also be compared at country and ecoregion levels, on the DOPA Explorer website: http://dopa-explorer.jrc.ec.europa.eu/dopa_explorer/.

Data updates Planned with each update of DOPA.

Codes Standard GIS operations applied to vector and raster data.

Methodology

Methodology The AGCI is based on the information provided by the global terrestrial biomass map developed by the GlobBiomass project, which estimates, with a spatial resolution of 100 m and for the reference year 2010, the amount of AGB (Mg/ha) considering the oven-dry weight woody parts (stem, bark, branches and twigs) of all living trees excluding stump and roots. The AGB values have been here converted to carbon content (AGCI) using the conversion factor of 0.5 (Mg C / Mg dry matter), which is consistent with the approach in the Good Practice Guidance in LULUCF by the IPCC (2003) and within the range of values (0.47 - 0.51) used in the related literature (Ruesch and Gibbs, 2008; Thurner *et al.*, 2014).




Figure 1. Screen capture of the above ground carbon content as presented in DOPA Explorer for a protected area, here the Matheniko wildlife reserve in Uganda.

The GlobBiomass map has been developed through a synergistic mapping approach and multiple estimation procedures combining spaceborne SAR (ALOS PALSAR, Envisat ASAR), optical (Landsat 7) and LiDAR (ICESAT) datasets, further supported by auxiliary products derived from earth observation (land cover, land surface temperature etc.) and in situ information. The earth observation data were used to estimate the Growing Stock Volume (GSV) of trees. GSV accounts for the volume of all living trees with more than 10 cm in diameter at breast height measured over bark from ground or stump height to a top stem diameter of 0 cm and excludes smaller branches, twigs, foliage, flowers, seeds, stump and roots (definition of FAO). Then, AGB was obtained from GSV with a

set of biomass expansion and conversion factors, derived from ground estimates of wood density and stem-to-total biomass expansion factors. See Quegan *et al.* (2017) for a detailed description of the algorithms and methods used in the production of the GlobBiomass map.

The AGB map data, with a spatial resolution of 100 m, were overlaid with the boundaries of each terrestrial or coastal protected area of size $\geq 25 \text{ km}^2$ to calculate the minimum, maximum, mean and standard deviation of AGCI ($\text{Mg C}/\text{km}^2$) within each protected area and the total AGC stored (Mg) in each protected area. UNESCO Biosphere Reserves were discarded as well as protected areas with known areas but undefined boundaries.

Input datasets The indicator uses the following input datasets:

Protected Areas

- WDPA of July 2018 (UNEP-WCMC & IUCN, 2018).
 - Latest version available from: www.protectedplanet.net

Above-Ground Biomass

- GlobBiomass global map of forest above-ground biomass (Santoro, 2018).
 - The global above-ground biomass map is available for download, in the form of $40^\circ \times 40^\circ$ tiles, at: <http://globbiomass.org/products/global-mapping/>

References

- Achard, F., Beuchle, R., Mayaux, P., Stibig, H. J., Bodart, C., Brink, A., ... & Lupi, A. (2014). Determination of tropical deforestation rates and related carbon losses from 1990 to 2010. *Global change biology*, 20(8): 2540-2554. <https://doi.org/10.1111/gcb.12605>
- Baccini, A. G. S. J., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., ... & Samanta, S. (2012). Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature climate change*, 2(3): 182. <https://doi.org/10.1038/nclimate1354>
- Baccini, A., Walker, W., Carvalho, L., Farina, M., Sulla-Menashe, D., & Houghton, R. A. (2017). Tropical forests are a net carbon source based on aboveground measurements of gain and loss. *Science*, 358(6360): 230-234. DOI: [10.1126/science.aam5962](https://doi.org/10.1126/science.aam5962)
- Gallaun, H., Zanchi, G., Nabuurs, G. J., Hengeveld, G., Schardt, M., & Verkerk, P. J. (2010). EU-wide maps of growing stock and above-ground biomass in forests based on remote sensing and field measurements. *Forest Ecology and Management*, 260(3): 252-261. <https://doi.org/10.1016/j.foreco.2009.10.011>
- IPCC. (2013). Good Practice Guidance for Land Use, Land-Use Change and Forestry. Intergovernmental Panel on Climate Change. IPCC National Greenhouse Gas Inventories Programme. Available [here](#).
- Quegan, S., Rauste, Y., Bouvet, A., Carreiras, J., Cartus, O., Carvalhais, N., LeToan, T., Mermoz, S., Santoro, M. (2017). D6 – Global biomass map algorithm

theoretical basis document. Prepared by the GlobBiomass project for the European Space Agency (ESA-ESRIN) in response to ESRIN/Contract No. 4000113100/14/I_NB. Available at <http://globbiomass.org/products/global-mapping/>

Land Cover CCI. (2017). Product User Guide Version 2.0 http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf

Rozendaal, D.M.A., Santoro, M., Schepaschenko, D., Avitabile, V., Herold, M. (2017). D17 – Validation Report of the GlobBiomass project. Prepared for European Space Agency (ESA-ESRIN). In response to ESRIN/Contract No. 4000113100/14/I_NB. Available [here](#).

Ruesch, A., and Holly K. Gibbs. 2008. New IPCC Tier-1 Global Biomass Carbon Map For the Year 2000. Available online from the Carbon Dioxide Information Analysis Center [<http://cdiac.ess-dive.lbl.gov>], Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T., Salas, W., ... & Petrova, S. (2011). Benchmark map of forest carbon stocks in tropical regions across three continents. *Proceedings of the National Academy of Sciences*, 108(24): 9899-9904. <https://doi.org/10.1073/pnas.1019576108>

Santoro, M. (2018). GlobBiomass - global datasets of forest biomass. PANGAEA, <https://doi.org/10.1594/PANGAEA.894711>

Santoro, M., Cartus, O., Mermoz, S., Bouvet, A., Le Toan, T., Carvalhais, N., Rozendaal, D., Herold, M., Avitabile, V., Quegan, S., Carreiras, J., Rauste, Y., Balzter, H., Schullius, C., Seifert, F.M. (2018). GlobBiomass global above-ground biomass and growing stock volume datasets, available on-line at <http://globbiomass.org/products/global-mapping>

Thurner, M., Beer, C., Santoro, M., Carvalhais, N., Wutzler, T., Schepaschenko, D., ... & Schullius, C. (2014). Carbon stock and density of northern boreal and temperate forests. *Global Ecology and Biogeography*, 23(3): 297-310. <https://doi.org/10.1111/geb.12125>

UNEP-WCMC & IUCN. (2018). Protected Planet: The World Database on Protected Areas (WDPA) [On-line], [July/2018], Cambridge, UK: UNEP-WCMC and IUCN. <http://www.protectedplanet.net>

Zarin, D. J., Harris, N. L., Baccini, A., Aksenov, D., Hansen, M. C., Azevedo-Ramos, C., ... & Allegretti, A. (2016). Can carbon emissions from tropical deforestation drop by 50% in 5 years? *Global change biology*, 22(4): 1336-1347. <https://doi.org/10.1111/gcb.13153>

Contact

Please contact us at: JRC-DOPA@ec.europa.eu

Factsheet last updated

March 27, 2019